

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 9/12	A1	(11) International Publication Number: WO 99/06025 (43) International Publication Date: 11 February 1999 (11.02.99)
(21) International Application Number: PCT/US98/16111 (22) International Filing Date: 4 August 1998 (04.08.98) (30) Priority Data: 08/904,565 4 August 1997 (04.08.97) US (71) Applicant: BAKER NORTON PHARMACEUTICALS, INC. [US/US]; 4400 Biscayne Boulevard, Miami, FL 33137 (US). (72) Inventor: AHMED, Tahir, 1020 South Greenway Drive, Coral Gables, FL 33134 (US). (74) Agent: SCHIFFMILLER, Martin, W.; Kirschstein, Ottinger, Israel & Schiffmiller, P.C., 489 Fifth Avenue, New York, NY 10017-6105 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: METHOD AND COMPOSITIONS FOR TREATING LATE PHASE ALLERGIC REACTIONS AND INFLAMMATORY DISEASES (57) Abstract A method of treating a mammalian patient suffering from or prone to a condition characterized by late phase allergic reactions, airway hyperresponsiveness or inflammatory reactions, e.g., asthma, allergic rhinitis, allergic dermatitis, allergic conjunctivitis, inflammatory bowel disease or rheumatoid arthritis, comprising the administration to the patient of an oral, parenteral, intrabronchial, topical, intranasal or intraocular pharmaceutical composition containing in each dose about 0.005 to about 1.0 mg per kilogram of patient body weight of ultra-low molecular weight heparins (ULMWH) or other sulfated polysaccharides having average molecular weights of about 1,000-3,000 daltons. Suitable inhalant and other pharmaceutical compositions for use in the novel treatment method are also disclosed.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

METHOD AND COMPOSITIONS FOR TREATING LATE PHASE ALLERGIC REACTIONS AND INFLAMMATORY DISEASES

BACKGROUND OF THE INVENTION

1. Field of the Invention

5 The invention relates to methods and compositions for preventing and reversing the symptoms and manifestations of late phase allergic reactions and inflammatory diseases.

2. Description of the Prior Art

10 Chronic asthma can be considered to be predominantly an inflammatory disease with associated bronchospasm. The degree of reactivity and narrowing of the bronchi in response to stimuli is greater in asthmatics than in normal individuals. Persistent inflammation is responsible for the bronchial hyperreactivity or airway hyperresponsiveness (AHR). Mucosal edema, mucus plugging and hypersecretion may be present; pulmonary parenchyma is normal. Airway narrowing may reverse
15 spontaneously or with therapy. Type 1 (immediate) immune responses may play an important role in the development of asthma in children and many adults; however, when onset of disease occurs in adulthood, allergic factors may be difficult to identify. Exposure to cold dry air, exercise and other aggravating factors also may trigger asthma.

20 The general goals of drug therapy for asthma are prevention of bronchospasm and long-term control of bronchial hyperreactivity. Because it is usually not possible for either patient or physician to predict when bronchospasm may

- 2 -

occur, patients with all but the most episodic and/or entirely seasonal attacks may require continuous therapy.

Beta agonists are useful as bronchodilator agents; they stimulate beta₂-adrenergic receptors, increase intracellular cAMP, and may inhibit the release of mast cell mediators. Other useful drugs include theophylline and related xanthine drugs, which produce bronchodilation through unknown mechanisms; the bismolone, cromolyn, which prevents the release of mediator substances and blocks respiratory neuronal reflexes; and corticosteroids, which primarily decrease inflammation and edema. Anticholinergic drugs may relieve bronchospasm by blocking parasympathetic cholinergic impulses at the receptor level. Antihistamines occasionally prevent or abort allergic asthmatic episodes, particularly in children, but they can only be partially effective in asthma because histamine is only one of many mediators.

The current drug modalities used for treatment of allergy-induced asthma suffer from a number of drawbacks. In general, the conventional agents have a relatively short duration of action and may be partially or wholly ineffective when administered after antigen challenge occurs. Moreover, because of serious adverse effects associated with the use of agents such as beta₂-adrenergic agonists and corticosteroids, the therapeutic margin of safety with such agents is relatively narrow and patients using them must be carefully monitored.

Bronchial hyperreactivity (or AHR) is a hallmark of asthma and is closely related to underlying airway inflammation. Worsening of asthma and airway inflammation is associated with increase in bronchial hyperreactivity, which can be induced by both antigenic and non-antigenic stimuli. Beta₂-adrenergic agonists are potent agents for the treatment of bronchospasm, but have no effect on airway

- 3 -

inflammation or bronchial hyperreactivity. In fact, chronic use of beta₂-adrenergic agents alone, by causing down regulation of beta₂-receptors, may worsen bronchial hyperreactivity. At present, corticosteroids are the only effective agents available which diminish bronchial hyperreactivity. Although inhaled corticosteroids are relatively safe in adult patients with asthma, these agents have tremendous toxicity in children, including adrenal suppression and reduced bone density and growth. Thus, the search for safer and effective agents which diminish bronchial hyperreactivity continues.

Patients with allergic asthma, following an inhalation challenge with the specific antigen exhibit at least two different patterns of bronchial responses. The majority of subjects develop an acute bronchoconstrictor response only, which resolves spontaneously within 1-3 hours; these subjects are termed "acute responders". A smaller number of subjects, however, develop both an early and a late response. These subjects are termed "dual responders". In dual responders, the acute response, which resolves spontaneously, is followed 4-12 hours later by a secondary increase in airway resistance ("late phase response"). Late responses and thus dual responders are of clinical importance, because of their association with prolonged airway hyperreactivity or hyperresponsiveness (AHR), worsening of symptoms and generally worse form of clinical asthma, requiring aggressive therapy.

Pharmacological studies in allergic animals have demonstrated that not only the bronchoconstrictor response but also the inflammatory cell influx and the mediator release pattern in dual responders is quite different from acute responders. Whereas histamine is the likely bronchoconstrictor mediator during acute phase, activated products of the lipoxygenase pathway (i.e., leukotrienes) may be the major

- 4 -

mediator involved in late phase reaction. Mast cells, however, have a central role in IgE-mediated allergic airway responses, and cromolyn sodium (a mast-cell membrane stabilizer), theoretically should prevent bronchoconstrictor responses in both "acute" and "dual" responders. Heterogeneity of mast cell subtypes may play a significant role in divergent responses and it may be dependent upon differences in signal transduction (second messenger system).

It has been discovered in the past several years that heparin administered intrabronchially can be an effective inhibitor of bronchospasm and bronchoconstriction and is consequently of value in the prophylaxis of asthma (see, e.g., Ahmed et al., New Eng. J. Med., 329:90-95, 1993; Ahmed, Resp. Drug Deliv., IV:55-63, 1994). It has been discovered further that low molecular weight heparins, e.g., heparins with an average molecular weight of 4,000-5,000 daltons, effectively prevent antigen-induced bronchoconstriction; these low molecular weight heparins also exhibit considerably less anticoagulant activity than commercial heparin, a desirable property when these agents are used in the treatment of asthma (see Ashkin et al., Am. Rev. Resp. Dis., 1993 Intl. Conf. Abstracts, p. A660). Both commercial and low-weight heparins are not effective, however, in suppressing AHR when administered after the patient has been exposed to antigen.

In parent application Serial No. 08/516,786 we disclosed that ultra-low molecular weight heparins (ULMWH) having an average molecular weight less than about 3,000 daltons are effective in suppressing AHR in acute asthmatic responders, even when administered after the patient has been challenged with antigen. However, experimental and clinical studies have shown that while inhaled commercial heparin can also attenuate early phase antigen-induced bronchoconstriction in acute

- 5 -

responders (though not after antigen challenge) it is ineffective in the treatment of dual responders. Hence, there was still considerable doubt after our earlier work with ULMWH as to whether these substances would show efficacy in the treatment of dual or late responders as they do in acute responders.

5 The current, conventional therapeutic modalities for asthmatic patients who are dual responders are generally a more aggressive and time-prolonged version of the therapies practiced on acute responders, described above. However, these therapies are not particularly effective in suppressing AHR, as noted previously, and, as a result of their generally short duration of action, cannot prevent the late phase
10 reaction and AHR observed in dual responders.

 It should be noted, however, that the airways are merely a prototype of organs or tissues affected by late phase reactions (LPR's). It has been established in the medical literature that the late phase bronchoconstriction and AHR observed in dual responder asthmatic patients is not an isolated phenomenon restricted to
15 asthmatic or even pulmonary conditions. There are cutaneous, nasal, ocular and systemic manifestations of LPR's in addition to the pulmonary ones. These allergic LPR phenomena are considered closely interrelated from the point of view of immunologic mechanisms. See Lemanske and Kaliner, "Late Phase Allergic Reactions", published in Allergies, Principles and Practice (Mosby Yearbook, Inc., 4th
20 ed. 1997). According to the latest understanding of LPR mechanisms it appears that the clinical diseases (whether of the skin, lung, nose, eye or other organs) recognized to involve allergic mechanisms have a histologic inflammatory component which follows the immediate allergic or hypersensitivity reaction that occurs on antigen challenge. This sequence of response appears to be connected to mast cell

- 6 -

m diators and propagated by other resident cells within target organs or by cells recruited into the sites of mast cell or basophilic degranulation. Corticosteroids which have proven valuable in the management of various allergic diseases and asthma may be beneficial because of their ability to attenuate this inflammatory process.

5 Furthermore, there are extra-pulmonary diseases where inflammatory response plays a major role, for example, inflammatory bowel disease, rheumatoid arthritis, glomerulonephritis and inflammatory skin disease. These conditions are also often treated with anti-inflammatory agents which may be of short duration or which, like steroidal and non-steroidal anti-inflammatory drugs, may frequently cause adverse
10 systemic or gastrointestinal reactions.

 Improved pharmaceutical treatments for late phase allergic reactions and inflammatory diseases are required.

SUMMARY OF THE INVENTION

 It is an object of this invention to provide more effective and safer methods
15 and compositions for treatment of conditions characterized by late phase allergic reactions or inflammatory reactions.

 It is another object of the present invention to provide a method and compositions for treatment of antigen-induced late phase asthma and bronchial hyperreactivity which do not suffer from the drawbacks of the prior art.

20 It is a further object of the present invention to provide a method and compositions for the treatment of asthmatic dual responders which are effective in preventing and reversing the manifestations of late phase asthma.

- 7 -

Still another object of the present invention is to provide a method and compositions as described above which are highly effective in diminishing specific and non-specific bronchial hyperreactivity, and even when administered after antigen challenge to the patient.

5 In keeping with these objects and others which will become apparent hereinafter, the invention resides in a method of treating a mammalian patient suffering from a condition which is characterized by late phase allergic reactions, including, e.g., pulmonary, nasal, cutaneous, ocular and systemic LPR's, or which is characterized by inflammatory reactions, through the intrabronchial, oral, topical,
10 parenteral, intranasal or intraocular administration to the patient of a pharmaceutical composition comprising from about 0.005 to about 1.0 mg of ultra-low molecular weight heparins (ULMWH) per kilogram of patient body weight in each dose. The administration of these heparins can be on an acute basis such as following antigen challenge, or on a chronic basis to suppress inflammatory reactions such as bronchial
15 hyperreactivity in asthma patients.

The ULMWH effective in the method of the invention have average molecular weights of about 1,000 to about 3,000 daltons and may exhibit a low level of anticoagulant activity or substantially no anticoagulant activity at all. Novel pharmaceutical compositions are also provided including, e.g., inhalant (intrabronchial)
20 compositions in the form of liquid or powder nebulizer or aerosol compositions containing suitable concentrations of these ULMWH.

- 8 -

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a graph illustrating the effects of antigen challenge on two groups of allergic sheep, one composed of acute responders and the other of dual responders. Data for each group are shown as antigen-induced mean \pm SE% change in SR_L (specific lung resistance), shown before (baseline), immediately after (P.A.) and up to 8 hours post-antigen.

+ = Significantly different from baseline ($P < .05$)

* = Significantly different from dual responders ($P < .05$)

FIGS. 2A and 2B comprise two graphs illustrating the differential effects of inhaled commercial heparin on antigen-induced bronchoconstriction in two groups of allergic sheep, one composed of acute responders ($n=8$) and the other of dual responders ($n=13$).

+ = Significantly different from baseline ($P < .05$)

* = Significantly different from control ($P < .05$)

FIG. 3 is a graph illustrating the effect of pretreatment with inhaled FragminTM (avg. mol. wt. 5,030 daltons) at 5.0 mg/kg on antigen-induced bronchoconstriction in dual responder allergic sheep. Data are shown as mean SR_L (in cm $H_2O/L/sec$) in a group of animals exposed to antigen, first with no drug treatment and again several days later after pretreatment with Fragmin.

FIG. 4 is a bar graph illustrating the effect of pretreatment with inhaled FragminTM at 5.0 mg/kg on AHR in allergic sheep. Data are shown as mean \pm SE PD_{400} in breath units at baseline and 24 hours post-antigen challenge in a group of animals exposed to antigen, first with no drug treatment and again several days later after pretreatment with Fragmin.

- 9 -

PD_{400} = Cumulative provoking dose of carbachol, increasing SR_L to 400% above baseline

+ = Significantly different from baseline ($P < .05$)

FIG. 5 is a graph illustrating the effect of pretreatment with inhaled
5 CY-216 (avg. mol. wt. 4,270 daltons) at 1.25 mg/kg on antigen-induced bronchoconstriction in dual responder allergic sheep. Data are shown as antigen-induced mean \pm SE% change in SR_L in a group of animals exposed to antigen, first with no drug treatment and again several days later after pretreatment with CY-216.

FIG. 6 is a bar graph illustrating the effect of pretreatment with inhaled CY-
10 216 at 1.25 mg/kg on AHR in allergic sheep. Data are shown as mean \pm SE PD_{400} in breath units at baseline and 24 hours post-antigen challenge in a group of animals exposed to antigen, first with no drug treatment and again several days later after pretreatment with CY-216.

+ = Significantly different from baseline ($P < .05$)

15 FIG. 7 is a graph illustrating the effect of pretreatment with inhaled ULMWH CY-222 (avg. mol. wt. 2,355 daltons) at 1.0 mg/kg on antigen-induced bronchoconstriction in dual responder allergic sheep. Data are shown as antigen-induced mean \pm SE% change in SR_L in a group of animals exposed to antigen, first with no drug treatment and again several days later after pretreatment with CY-222.

20 * = Significantly different from control ($P < .05$)

FIG. 8 is a bar graph illustrating the effect of pretreatment with inhaled CY-222 at 1.0 mg/kg on AHR in allergic sheep. Data are shown as mean \pm SE PD_{400} in breath units at baseline and 24 hours post-antigen challenge in a group of animals

- 10 -

exposed to antigen, first with no drug treatment and again several days later after pretreatment with CY-222.

+ = Significantly different from baseline ($P < .05$)

FIG. 9 is a graph illustrating the effect of treatment post-antigen challenge with inhaled CY-222 at 1.0 mg/kg on antigen-induced bronchoconstriction in dual responder allergic sheep. Data are shown as antigen-induced mean \pm SE% change in SR_L , shown before, immediately after (time zero) and up to 8 hours post-antigen, in a group of animals exposed to antigen, first with no drug treatment and again several days later when CY-222 was administered immediately after the post-antigen measurement of SR_L (arrow).

* = Significantly different from control ($P < .05$)

FIG. 10 is a bar graph illustrating the effect of treatment post-antigen challenge (arrow in FIG. 9) with inhaled CY-222 at 1.0 mg/kg on AHR in allergic sheep. Data are shown as mean \pm SE PD_{400} in breath units at baseline and 24 hours post-antigen challenge in a group of animals exposed to antigen, first with no drug treatment and again several days later when CY-222 was administered immediately after antigen challenge.

+ = Significantly different from baseline ($P < .05$)

FIG. 11 is a graph illustrating the effect of pretreatment with inhaled ULMWH FRU-70 (avg. mol. wt. 2,500 daltons) at 1.0 mg/kg on antigen-induced bronchoconstriction in dual responder allergic sheep. Data are shown as antigen-induced mean \pm SE% change in SR_L in a group of animals exposed to antigen, first with no drug treatment and again several days later after pretreatment with FRU-70.

* = Significantly different from control ($P < .05$)

- 11 -

FIG. 12 is a bar graph illustrating the effect of pretreatment with inhaled FRU-70 at 1.0 mg/kg on AHR in allergic sheep. Data are shown as mean \pm SE PD₄₀₀ in breath units at baseline and 24 hours post-antigen challenge in a group of animals exposed to antigen, first with no drug treatment and again several days later after pretreatment with FRU-70.

+ = Significantly different from baseline (P<.05)

FIG. 13 is a graph illustrating the effect of treatment post-antigen challenge with inhaled FRU-70 at 0.5 mg/kg on antigen-induced bronchoconstriction in dual responder allergic sheep. Data are shown as antigen-induced mean \pm SE% change in SR_L, shown before, immediately after (time zero) and up to 8 hours post-antigen, in a group of animals exposed to antigen, first with no drug treatment and again several days later when FRU-70 was administered immediately after the post-antigen measurement of SR_L (arrow).

* = Significantly different from control (P<.05)

FIG. 14 is a bar graph illustrating the effect of treatment post-antigen challenge (arrow in FIG. 13) with inhaled FRU-70 at 0.5 mg/kg on AHR in allergic sheep. Data are shown as mean \pm SE PD₄₀₀ in breath units at baseline and 24 hours post-antigen challenge in a group of animals exposed to antigen, first with no drug treatment and again several days later when FRU-70 was administered immediately after antigen challenge.

+ = Significantly different from baseline (P<.05)

FIG. 15 is a graph illustrating the effect of treatment post-antigen challenge with inhaled hexasaccharide mixture (avg. mol. wt. 1,930 daltons) at 0.5 mg/kg on antigen-induced bronchoconstriction in dual responder allergic sheep. Data are

- 12 -

shown as antigen-induced mean \pm SE% change in SR_L , shown before, immediately after (time zero) and up to 8 hours post-antigen, in a group of animals exposed to antigen, first with no drug treatment and again several days later when the hexasaccharide mixture was administered immediately after the post-antigen measurement of SR_L (arrow).

* = Significantly different from control ($P < .05$)

FIG. 16 is a bar graph illustrating the effect of treatment post-antigen challenge (arrow in FIG. 15) with inhaled hexasaccharide mixture at 0.5 mg/kg on AHR in allergic sheep. Data are shown as mean \pm SE PD_{400} in breath units at baseline and 24 hours post-antigen challenge in a group of animals exposed to antigen, first with no drug treatment and again several days later when the hexasaccharide mixture was administered immediately after antigen challenge.

+ = Significantly different from baseline ($P < .05$)

FIG. 17 is a graph illustrating the effect of treatment post-antigen challenge with inhaled purified hexasaccharide (avg. mol. wt. 1998 daltons) at 0.062 mg/kg on antigen-induced bronchoconstriction in dual responder allergic sheep. Data are shown as antigen-induced mean \pm SE% change in SR_L , shown before, immediately after (time zero) and up to 8 hours post-antigen, in a group of animals exposed to antigen, first with no drug treatment and again several days later when the hexasaccharide was administered immediately after the post-antigen measurement of SR_L (arrow).

* = Significantly different from control ($P < .05$)

FIG. 18 is a bar graph illustrating the effect of treatment post-antigen challenge (arrow in FIG. 17) with inhaled purified hexasaccharide at 0.062 mg/kg on

- 13 -

AHR in allergic sh ep. Data are shown as mean \pm SE PD₄₀₀ in breath units at baseline and 24 hours post-antigen challenge first, with no drug treatment and again several days later when the hexasaccharide was administered immediately after antigen challenge.

5 + = Significantly different from baseline (P<.05)

FIG. 19 is a graph illustrating the effect of treatment post-antigen challenge with inhaled purified tetrasaccharide (avg. mol. wt. 1290 daltons) at 0.062 mg/kg on antigen-induced bronchoconstriction in dual responder allergic sheep. Data are shown as antigen-induced mean \pm SE% change in SR_L, shown before, immediately
10 after (time zero) and up to 8 hours post-antigen, in a group of animals exposed to antigen, first with no drug treatment and again several days later when the tetrasaccharide was administered immediately after the post-antigen measurement of SR_L (arrow).

* = Significantly different from control (P<.05)

15 FIG. 20 is a bar graph illustrating the effect of treatment post-antigen challenge (arrow in FIG. 19) with inhaled purified tetrasaccharide at 0.062 mg/kg on AHR in allergic sheep. Data are shown as mean \pm SE PD₄₀₀ in breath units at baseline and 24 hours post-antigen challenge first, with no drug treatment and again several days later when the tetrasaccharide was administered immediately after
20 antigen challenge. + = Significantly different from baseline (P<.05)

FIG. 21 is a graph illustrating the effect of treatment post-antigen challenge with inhaled octasaccharide (avg. mol. wt. 2,480 daltons) at 0.25 mg/kg on antigen-induced bronchoconstriction in dual responder allergic sheep. Data are shown as antigen-induced mean \pm SE% change in SR_L, shown before, immediately after (time

- 14 -

zero) and up to 8 hours post-antigen, in a group of animals exposed to antigen, first with no drug treatment and again several days later when the octasaccharide mixture was administered immediately after the post-antigen measurement of SR_L (arrow).

* = Significantly different from control ($P < .05$)

5 FIG. 22 is a bar graph illustrating the effect of treatment post-antigen challenge (arrow in FIG. 21) with inhaled octasaccharide at 0.25 mg/kg on AHR in allergic sheep. Data are shown as mean \pm SE PD_{400} in breath units at baseline and 24 hours post-antigen challenge in a group of animals exposed to antigen, first with no drug treatment and again several days later when the octasaccharide was
10 administered immediately after antigen challenge.

+ = Significantly different from baseline ($P < .05$)

 FIG. 23 is a graph illustrating the effect of pretreatment with inhaled disaccharide (avg. mol. wt. 660 daltons) at 1.0 mg/kg on antigen-induced bronchoconstriction in dual responder allergic sheep. Data are shown as antigen-
15 induced mean \pm SE% change in SR_L , shown before, immediately after (time zero) and up to 8 hours post-antigen, in a group of animals exposed to antigen, first with no drug treatment and again several days later after pretreatment with disaccharide.

 FIG. 24 is a graph illustrating the effect of pretreatment with orally administered purified hexasaccharide (avg. mol. wt. 1,998 daltons) at 2.0 mg/kg on
20 antigen-induced bronchoconstriction in a dual responder allergic sheep. Data are shown as antigen-induced mean \pm SE% change in SR_L in a single sheep exposed to antigen, first with no drug treatment and again several days later after pretreatment with hexasaccharide.

- 15 -

FIG. 25 is a bar graph illustrating the effect of pretreatment with orally administered purified hexasaccharide at 2.0 mg/kg on AHR in an allergic sheep. Data are shown as mean \pm SE PD₄₀₀ in breath units at baseline and 24 hours post-antigen challenge in a single sheep exposed to antigen, first with no drug treatment and again several days later after pretreatment with hexasaccharide.

FIG. 26 is a graph illustrating the effect of pretreatment with intravenously administered purified hexasaccharide (avg. mol. wt. 1,998 daltons) at 0.25 mg/kg on antigen-induced bronchoconstriction in a dual responder allergic sheep. Data are shown as antigen-induced mean \pm SE% change in SR_L in a single sheep exposed to antigen, first with no drug treatment and again several days later after pretreatment with hexasaccharide.

FIG. 27 is a bar graph illustrating the effect of pretreatment with intravenously administered purified hexasaccharides at 0.25 mg/kg on AHR in allergic sheep. Data are shown as mean \pm SE PD₄₀₀ in breath units at baseline and 24 hours post-antigen challenge in a single sheep exposed to antigen, first with no drug treatment and again several days later after pretreatment with hexasaccharide.

FIG. 28 is a bar graph illustrating the comparative activity in preventing antigen-induced eosinophil influx in the bronchoalveolar lavage fluid of three groups of mice administered, respectively, inhaled, oral and intraperitoneal purified hexasaccharide.

DETAILED DESCRIPTION OF THE INVENTION

The present invention pertains generally to a method of treatment of mammalian patients suffering from or prone to development of disease conditions

- 20 -

While the sulfated polysaccharides used in the method and compositions of the invention are generally referred to herein as ultra-low molecular weight heparins, i.e., ultra-low molecular weight fractions derived from naturally occurring heparin (or synthetic versions of such ULMWH), the invention may also encompass the use of sulfated polysaccharides derived from heparan sulfate, dermatan sulfate, chondroitin sulfate, pentosan polysulfate and/or other glycosaminoglycans and mucopolysaccharides. The subject sulfated polysaccharide fractions must, however, have an average molecular weight of about 1,000-3,000 daltons. Pharmaceutically acceptable salts of the effective ULMWH or any of the other sulfated polysaccharides listed above may also be utilized, e.g., the sodium, calcium or potassium salts.

In accordance with this first aspect of the invention, a human or other mammalian patient who is a dual responder who has inhaled, ingested or otherwise come into contact with an antigen (i.e., has been "challenged" with an antigen) of a type known to provoke asthmatic episodes in that patient, or a patient who may be exposed at a future time to antigen challenge, is administered via inhalation at least one dose of a pharmaceutical composition containing one or more effective ULMWH cumulatively present in the above-described concentration ranges. Additional doses may subsequently be administered as necessary after antigen challenge until the patient regains or maintains normal airflow resistance levels.

The invention also comprehends in a second aspect the chronic administration of effective ULMWH to dual responder asthma patients to reduce and suppress early and late phase AHR. "Chronic administration" as used herein refers to administration of pharmaceutical compositions containing effective ULMWH at least once daily for at least ten consecutive days. Chronic administration of a composition containing from about 0.005-1.0 mg/kg per dose, and preferably about 0.0075-0.75

- 21 -

mg/kg per dose, can be continued indefinitely to provide AHR-suppressant therapy at least comparable to corticosteroids but substantially without side effects.

The inhalant (intrabronchial) ULMWH compositions used in the present invention to treat late phase asthma and other pulmonary conditions may comprise liquid or powder compositions containing effective ULMWH fractions and suitable for nebulization and intrabronchial use, or aerosol compositions administered via an aerosol unit dispensing metered doses.

Suitable liquid compositions comprise for example, effective ULMWH in an aqueous, pharmaceutically acceptable inhalant solvent, e.g., isotonic saline or bacteriostatic water. The solutions are administered by means of a pump or squeeze-actuated nebulized spray dispenser, or by any other conventional means for causing or enabling the requisite dosage amount of the liquid composition to be inhaled into the mammalian patient's lungs.

Suitable powder compositions include, by way of illustration, powdered preparations of heparin thoroughly intermixed with lactose or other inert powders acceptable for intrabronchial administration. The powder compositions can be administered via an aerosol dispenser or encased in a breakable capsule which may be inserted by the mammalian patient into a device that punctures the capsule and blows the powder out in a steady stream suitable for inhalation.

Aerosol formulations for use in the subject method would typically include fluorinated alkane propellants, surfactants and co-solvents and may be filled into aluminum or other conventional aerosol containers which are then closed by a suitable metering valve and pressurized with propellant, producing a metered dose inhaler (MDI).

- 22 -

The total concentration of effective ULMWH in any propellant vehicle suitable for use in a pressured aerosol dispenser, such as an MDI, must be sufficiently high to provide a dose of about 0.005-0.1 mg (5-100 μ g) of effective ULMWH per kilogram of patient body weight per administration. Thus, for example, if an MDI
5 delivers about 85 μ l of drug-containing propellant vehicle per actuation, the concentration of effective ULMWH in the vehicle in the case of a mammalian patient weighing 75 kg would be approximately 0.0045-0.088 mg/ μ l (4.5-88 μ g/ μ l), delivering 0.375 to 7.5 mg (375-7,500 μ g) of ULMWH per actuation, if it is desired to deliver the entire dose with a single actuation. If a two-actuation dose is desired, the
10 corresponding concentration range would be approximately 0.0022-0.044 mg/ μ l (2.2-44 μ g/ μ l), delivering 0.188 to 3.75 mg (188-3,750 μ g) of ULMWH per actuation.

The total concentration of effective ULMWH in any liquid nebulizer solution must be sufficiently high to provide a dose of about 0.05-1.0 mg (50-1000 μ g) of effective ULMWH per kilogram of patient body weight per administration. Thus, for
15 example, if the nebulizer utilized delivers 5 ml of solution per actuation, the concentration of effective ULMWH in the case of a mammalian patient weighing 75 kg should be approximately 0.75-15.0 mg/ml.

In a further aspect of the invention, effective ULMWH-containing compositions are administered orally or parenterally (e.g., IV or IM) to mammalian
20 patients suffering from antigen-induced late phase asthma, i.e., who are dual responders, prior to exposure of the patient to antigen-challenge. The oral or parenteral compositions contain about 0.005 to about 1.0 mg of effective ULMWH per kg of patient body weight in each dose. The oral or parenteral compositions may be administered up to 8 hours (but preferably not more than 4 hours) prior to antigen

- 23 -

challenge and are effective in reducing early and late phase bronchoconstriction and in suppressing AHR.

As those skilled in the pharmaceutical arts will appreciate, many conventional methods and apparatus are available for administering precisely metered doses of intrabronchial medicaments and for regulating the desired dosage amount in accordance with patient weight and the severity of the patient's condition. Moreover, there are many art-recognized liquid, powder and aerosol vehicles suitable for the intrabronchial ULMWH compositions of the present invention, and many pharmaceutically acceptable oral and parenteral vehicles which may be employed for the oral and parenteral ULMWH-containing compositions. The invention is not limited to any particular inert vehicles, solvents, carriers excipients or dosage forms and is not restricted to any particular methods or apparatus of intrabronchial administration.

The pharmaceutical compositions may also be dosage forms which contain the effective ULMWH as active ingredients in any pharmaceutically acceptable oral, injectable or IV dosage vehicles, or in topical or intraocular vehicles. Each dosage form includes about 0.005-1.0 mg/kg of average patient body weight of effective ULMWH (one or a combination of ULMWH) and pharmaceutically acceptable inert ingredients, e.g., conventional excipients, vehicles, fillers, binders, disintegrants, solvents, solubilizing agents, sweeteners, coloring agents and any other inactive ingredients which are regularly included in pharmaceutical dosage forms for oral administration. Suitable oral dosage forms include tablets, capsules, caplets, gelcaps, pills, liquid solutions, suspensions or elixirs, powders, lozenges, micronized particles and osmotic delivery systems. Injectable and IV dosage forms include isotonic saline solutions or dextrose solutions containing suitable buffers and preservatives. Many suitable dosage forms and vehicles, and listings of inactive

- 24 -

ingredi nts therefor, are well-known in the art and are set forth in standard texts such as Remington's Pharmaceutical Sciences, 17th edition (1985).

5 The ULMWH compositions described herein provide highly effective treatment for early and late phase antigen-induced asthma even after antigen challenge has occurred, as well as for other conditions characterized by late phase allergic reactions.

To demonstrate the unexpected superiority of the effective ULMWH in comparison with higher molecular weight heparins in treating asthmatic dual responders, experiments were conducted comparing the effects of different heparin
10 types on dual responder allergic sheep, both before and after antigen challenge. Detailed descriptions of these experiments and of the results obtained are set forth in the following examples as well as in the graphs shown in the drawings.

The following examples, while illustrating the methods and compositions of the invention and demonstrating the efficacy of the same, are not intended to set
15 forth specific compositions, materials, procedures or dosage regimens which must be utilized exclusively in order to practice the invention.

- 25 -

EXAMPLE I**Administration of Inhaled ULMWH to Dual Responder Allergic Sheep****Methods****Pulmonary Airflow Resistance:**

5 Allergic sheep with previously documented dual bronchoconstrictor response to Ascaris suum antigen were used for all studies. The sheep were intubated with a cuffed nasotracheal tube and pulmonary airflow resistance (R_L) was measured by the esophageal balloon catheter technique, while thoracic gas volume was measured by body plethysmography. Data were expressed as specific R_L (SR_L ,
10 defined as R_L times thoracic gas volume (V_{tg})).

Airway Responsiveness:

To assess airway responsiveness, cumulative dose-response curves to inhaled carbachol were performed by measuring SR_L before and after inhalation of buffered saline and after each administration of 10 breaths of increasing
15 concentrations of carbachol (0.25, 0.5, 1.0, 2.0 and 4.0% wt/vol solution). Airway responsiveness was measured by determining the cumulative provocation dose (PD_{400}) of carbachol (in breath units) that increased SR_L to 400% above baseline. One breath unit was defined as one breath of 1% carbachol solution.

Heparin Fractions:

20 In the studies reported herein various heparin materials were administered to the dual responder allergic sheep prior to and/or after antigen challenge. Some of these heparins were ULMWH fractions of average molecular weight between 1,000 and 3,000 daltons, some were of higher average molecular weight, and one was of lower molecular weight. The heparin fractions tested are set forth in Table 1 below.

TABLE 1**HEPARIN FRACTIONS AND THEIR MOLECULAR WEIGHTS**

	FRACTION	AVG. MOL. WT.	CATEGORY
	Commercial heparin	15,000 d	Heparin
5	Fragmin™	5,030 d	Low mol. wt.
	CY-216	4,270 d	Low mol. wt.
	CY-222 ¹ (Sanofi)	2,355 d	ULMWH
	FRU-70 ² (Kabivitrum)	2,500 d	ULMWH
	Hexasaccharide mixture ³	1,930 d	ULMWH
10	Octasaccharide ⁴	2,480 d	ULMWH
	Purified hexasaccharide ⁵	1,998 d	ULMWH
	Purified tetrasaccharide ⁶	1,290 d	ULMWH
	Disaccharide ⁷	660 d	Sub-ULMWH

¹ An anticoagulant octasaccharide mixture.

² A non-anticoagulant octasaccharide mixture.

³ Oligosaccharide derived from commercial porcine heparin, comprising primarily tetrasaccharide, hexasaccharide, octasaccharide and decasaccharide fractions.

⁴ Obtained from the hexasaccharide mixture by gel column chromatography, comprises about 70% octasaccharide and 30% decasaccharide fractions.

⁵ Obtained from the hexasaccharide mixture by gel column chromatography.

⁶ Obtained from the hexasaccharide mixture by gel column chromatography.

⁷ The disaccharide was trisulfated but had a molecular weight so low it could not be considered a ULMWH fraction with heparin-like properties.

- 27 -

Experimental Protocol

Airway Studies

Each animal's baseline airway responsiveness (PD_{400}) was determined, and then on different experimental days the sheep underwent airway challenge with Ascaris suum antigen. SR_L was measured, before and immediately after challenge, and then hourly for 8 hours. The post-challenge PD_{400} was measured 24 hours after antigen challenge when AHR occurred. The protocol was repeated at least 14 days later, but each animal was administered a dose of one of the test heparin fractions either about 30 minutes before antigen challenge or immediately after post-challenge SR_L measurement.

Data Analysis

Data were expressed as:

$$(a) \text{ } SR_L (\% \text{ change}) = \frac{\text{post challenge } SR_L - \text{baseline } SR_L \times 100}{\text{baseline } SR_L}$$

(b) PD_{400} (in breath units)

Results

FIG. 1 illustrates the differential reactions to antigen challenge of two groups of allergic sheep, one composed of acute responders and the other of dual responders. The SR_L of the acute responders returned to approximately baseline levels after about three hours post-antigen and remained there. In the dual responders, however, there is a late phase peak in SR_L at about six hours with levels remaining significantly above baseline through the eight-hour end point of the study. It is this second, late phase peak which characterizes dual responders.

- 28 -

FIGS. 2A and 2B depict the effects of pre-challenge treatment with inhaled commercial heparin on SR_L in acute responders (FIG. 2A) and in dual responders (FIG. 2B). While the SR_L of the acute responders remained at near-baseline levels even after antigen challenge, both the early phase and the late phase SR_L in the dual responders was not alleviated by heparin pretreatment, even in dual responders administered as much as 2000 units per kilogram.

FIGS. 3-6 illustrate the lack of efficacy of the low molecular weight heparin fractions, Fragmin and CY-216, in modifying either bronchoconstriction or AHR in dual responders when administered prior to antigen challenge.

FIGS. 7-10 show that both pretreatment and post-antigen challenge treatment with the inhaled ULMWH CY-222 (avg. mol. wt. 2355 d, within the range of effective ULMWH in accordance with the invention) were effective in significantly modifying both early and late phase antigen-induced bronchoconstriction and AHR in the dual responders.

FIGS. 11-14 illustrate the efficacy of both pretreatment and post-antigen challenge treatment with the ULMWH FRU-70 (avg. mol. wt. 2500 d) in the treatment of early and late phase asthma.

FIGS. 15-22 demonstrate the efficacy of various effective ULMWH fractions, even when administered post-antigen challenge, in significantly reducing bronchoconstriction and AHR in dual responders.

FIG. 23 shows that the disaccharide fraction, having an average molecular weight of only about 660 d (substantially below the weight range required for the effective ULMWH fractions) was ineffective in modifying antigen-induced bronchoconstriction in the dual responder allergic sheep.

- 29 -

In the experiments whose data are reflected in FIGS. 7-14 and 17-22 the dosage of the effective ULMWH administered to the allergic sheep was the lowest effective dose (determined through dose-ranging trials) for each ULMWH fraction. It will be observed that the different ULMWH had varying minimum effective dose levels in the treatment of dual responders. The minimum effective dose was about 1.0 mg/kg for CY-222 and for FRU-70 administered prior to antigen challenge, but about 0.5 mg/kg for FRU-70 administered post-antigen challenge and for the inhaled hexasaccharide mixture. The purified tetrasaccharide, having the lowest average molecular weight of any of the effective ULMWH tested, had a minimum effective dose when administered post-antigen of 0.062 mg/kg, as did the purified hexasaccharide. These data tend to suggest that purified fractions having an average weight near the lower limit of about 1,000 d may be the most effective ULMWH, at least in the treatment of dual responders. The optimal structural domain and/or sequence for the observed antiallergic and/or anti-inflammatory activity appears to be the tetrasaccharide.

- 30 -

EXAMPLE 2**Administration of Oral ULMWH to Dual Responder Allergic Sheep**

The procedure of Example 1, in terms of the test animals and evaluation methods, were followed in the present experiment.

5 One dual-responder allergic sheep was administered orally 2 mg/kg of purified hexasaccharide (avg. mol. wt. 1998 daltons) 90 minutes before challenge with Ascaris Suum antigen. The effects of the pretreatment with the hexasaccharide on SR_L from baseline (time of administration of the hexasaccharide) through 8 hours after antigen challenge is reflected on FIG. 24. Also shown on FIG. 24 for comparison
10 purposes is the percentage change in SR_L in the same dual responder sheep (in an experiment conducted several days earlier) challenged with antigen but without ULMWH pretreatment.

Shown in FIG. 25 are the respective PD_{400} values measured at baseline and post-antigen when the sheep was administered antigen challenge with
15 hexasaccharide pretreatment and with no pretreatment (control).

EXAMPLE 3**Administration of Intravenous ULMWH to Dual Responder Allergic Sheep**

The procedure of Example 2 was followed with another dual responder allergic sheep, except that 0.25 mg/kg of purified hexasaccharide was administered
20 intravenously one hour before antigen challenge in one experiment, while antigen was administered with no pretreatment in the second (control) experiment. The percentage change in SR_L for the pretreatment and control experiments are shown in

- 31 -

FIG. 26 and the PD_{400} values for those experiments at baseline and post-antigen are shown in FIG. 27.

EXAMPLE 4

Prevention of Antigen-induced Eosinophil Influx in Mice

5 In four groups of sensitized laboratory mice (n=3 in each group) bronchoalveolar lavage was performed 24 hours after antigen challenge to determine eosinophil influx values in each group. The mice were treated with either aerosolized saline (placebo) or purified hexasaccharide administered by the following routes and in the following dosage amounts, respectively: inhaled aerosol,⁸ oral (100 µg) and
10 intraperitoneal (40 µg). The percentage inhibition of eosinophil influx effected in each treatment group was determined by comparing the level of such influx measured in bronchoalveolar lavage fluid subsequent to hexasaccharide administration with the saline group.

 The mean percentage inhibition values for the three treatment groups of mice
15 are reflected in FIG. 28. The mice receiving inhaled and oral hexasaccharide showed a 40-50% reduction in eosinophil influx while the mice receiving intraperitoneal hexasaccharide showed about a 20% reduction in such influx.

⁸ The mice (n=3) were placed in a chamber containing 10 mg of hexasaccharide in 9 ml of bacteriostatic injection water, which was aerosolized. The mice were allowed to inhale the aerosol for about 30 minutes.

- 32 -

The differential effects of commercial heparin observed in acute and dual responders (shown in FIGS. 2A and 2B) might indicate the involvement of different signaling pathways during airway anaphylaxis. This would suggest that during immunologically mediated mast-cell reaction in the airways, IP_3 is the predominantly active pathway in "acute responders" while non- IP_3 pathways (e.g., diacyl-glycerol/protein kinase C or other pathways) may be operative in "dual responders".

The late phase response and AHR are associated with marked airway inflammation. The pathological studies of the airway mucosa and bronchoalveolar lavage (BAL) have shown influx of eosinophils, neutrophils and activated T-lymphocytes during this phase. Increased levels of eosinophil-derived inflammatory mediators in plasma and BAL, including eosinophilic cationic protein and major basic protein, have been observed during the late phase reaction. Upregulation of TH_2 -type cytokines (IL_4 and IL_5) following allergen challenge has also been observed during the late phase. Thus, the cellular inflammatory response, in combination with released pro-inflammatory mediators (e.g., leukotrienes, PAF, eosinophilic proteins, etc.) and locally produced cytokines in the bronchial mucosa, play a central role in the late phase allergic inflammation and bronchoconstriction.

AHR and thus airway inflammation can be modified either by prevention of mast mediator release by "anti-allergic agents" (e.g., cromolyn sodium) or by the action of "anti-inflammatory" agents like glucocorticosteroids. The "anti-allergic" agents are only effective as prophylactic agents and can prevent the mediator release and AHR. Because these agents do not possess anti-inflammatory activity, they are generally ineffective when administered after the exposure to antigen. By contrast "anti-

- 33 -

inflammatory" agents can attenuate post-antigen AHR and airway inflammation, whether administered before or after the exposure to antigen. Our data suggest that the actions of ULMWH are analogous to the anti-inflammatory actions of glucocorticosteroids.

Since the effective ULMWH can modify AHR even when administered after antigen challenge, they should also be useful in the treatment of non-asthmatic conditions associated with AHR, e.g., chronic bronchitis, emphysema and cystic fibrosis.

Moreover, in view of our findings regarding the efficacy of certain ULMWH in inhibiting asthmatic LPR in a manner resembling the anti-inflammatory effects of corticosteroids, the effective ULMWH should be useful in the treatment of the following conditions and by the following routes of administration:

1. Late phase reactions and inflammatory response in extra-pulmonary sites:

- (a) allergic rhinitis
- (b) allergic dermatitis
- (c) allergic conjunctivitis

2. Extra-pulmonary diseases where inflammatory response plays a major role:

- (i) inflammatory bowel disease
- (ii) rheumatoid arthritis and other collagen vascular diseases
- (iii) glomerulonephritis
- (iv) inflammatory skin diseases
- (v) sarcoidosis

3. Routes of Administration

- (i) intrabronchial

- 34 -

(ii) intranasal

(iii) intraocular

(iv) topical

(v) oral

5 (vi) parenteral (IM or IV)

It should be emphasized, however, that the invention is not restricted or limited in any way to any theoretical or actual physiological or biochemical mechanisms or pathways, but comprehends the methods of treatment of conditions characterized by late phase allergic reactions, or treatment of dual responder mammalian patients, and the
10 compositions for use in said methods described hereinabove, notwithstanding the actual mechanisms of action involved.

It has thus been shown that there have been provided methods and compositions which achieve the various objects of the invention and which are well adapted to meet the conditions of practical use.

15 As various possible embodiments might be made of the above invention, and as various changes might be made in the embodiments set forth above, it is to be understood that all matters herein described are to be interpreted as illustrative and not in a limiting sense.

What is claimed as new and desired to be protected by Letters Patent is set forth in the following claims.

- 35 -

CLAIMS:

1. A method of treating a mammalian patient suffering from or prone to a condition characterized by late phase allergic reactions, airway hyperresponsiveness or inflammatory reactions, said method comprising the administration to the patient of a pharmaceutical composition containing about 0.005 to about 1.0 mg of ultra-low molecular weight heparins (ULMWH) per kilogram of patient body weight in each dose, said ULMWH having an average molecular weight of about 1,000 to about 3,000 daltons.

2. A method according to claim 1 wherein said ULMWH have an average molecular weight of about 1,000 to about 2,500 daltons.

3. A method according to claim 1 wherein said ULMWH comprise heparin fractions selected from the group consisting of tetrasaccharides, pentasaccharides, hexasaccharides, septasaccharides, octasaccharides and decasaccharides and pharmaceutically acceptable salts thereof.

4. A method according to claim 1 wherein said ULMWH are N-sulfated.

5. A method according to claim 1 wherein said composition contains about 0.075 to about 0.75 mg of said ULMWH per kilogram per dose.

6. A method according to claim 1 wherein said ULMWH have substantially no anticoagulant activity.

7. A method according to claim 1 wherein said composition is administered by the oral, parenteral, topical, intrabronchial, intranasal or intraocular routes.

8. A method according to claim 7 wherein said parenteral administration is intravenous or intramuscular.

- 36 -

9. A method according to claim 1 wherein said condition is characterized by late phase allergic reactions selected from the group consisting of pulmonary late phase reactions, nasal late phase reactions, cutaneous late phase reactions, ocular late phase reactions and systemic late phase reactions

10. A method according to claim 9 wherein said late phase reactions are pulmonary late phase reactions.

11. A method according to claim 10 wherein said condition is late phase asthma.

12. A method according to claim 1 wherein said condition is a non-asthmatic condition characterized by airway hyperresponsiveness.

13. A method according to claim 12 wherein said condition is selected from the group consisting of chronic bronchitis, emphysema and cystic fibrosis.

14. A method according to claim 1 wherein said condition is characterized by inflammatory reactions.

15. A method according to claim 1 wherein said condition is selected from the group consisting of allergic rhinitis, allergic dermatitis, allergic conjunctivitis, inflammatory bowel disease, rheumatoid arthritis, collagen vascular diseases, glomerulonephritis, inflammatory skin diseases and sarcoidosis.

16. A method according to claim 1 wherein said composition comprises a solution or suspension of said ULMWH in an aqueous, pharmaceutically acceptable, liquid inhalant vehicle.

17. A method according to claim 16 wherein said vehicle is isotonic saline or bacteriostatic water.

- 37 -

18. A method according to claim 16 wherein said composition is administered by means of a pump or squeeze-actuated nebulizer.

19. A method according to claim 16 wherein a sufficient amount of said composition is administered to the patient to provide a dose of about 0.05-1.0 mg/kg of said ULMWH.

20. A method according to claim 16 wherein said composition contains about 0.75-15.0 mg/ml of said ULMWH.

21. A method according to claim 1 wherein said composition is an aerosol composition comprising an aerosol propellant.

22. A method according to claim 21 wherein said composition is dispensed via a metered dose inhaler.

23. A method according to claim 21 wherein a sufficient amount of said composition is administered to the patient to provide a dose of about 0.005-0.1 mg/kg of said ULMWH.

24. A method according to claim 21 wherein said composition contains about 2.2-88 µg/µl of said ULMWH.

25. A method according to claim 1 wherein said composition comprises a powdered preparation of said ULMWH intermixed with an inert powder acceptable for intrabronchial administration.

26. A method according to claim 25 wherein said inert powder is lactose.

27. A method according to claim 25 wherein said composition is administered via an aerosol dispenser.

- 38 -

28. A method according to claim 25 wherein said composition is administered from a breakable capsule.

29. A method according claim 1 wherein said composition is administered to the patient prior to challenge with allergic reaction-inducing antigen.

30. A method according claim 1 wherein said composition is administered to the patient subsequent to challenge with allergic reaction-inducing antigen.

31. A pharmaceutical composition for the treatment of a mammalian patient suffering from or prone to a condition characterized by late phase allergic reactions, airway hyperresponsiveness or inflammatory reactions, said composition comprising about 0.005 to about 1.0 mg of ULMWH per kilogram of patient body weight in each dose in a pharmaceutically acceptable inhalant, oral, parenteral, topical, intranasal or intraocular vehicle, said ULMWH having an average molecular weight of about 1,000 to about 3,000 daltons.

32. A composition according to claim 31 wherein said ULMWH have an average molecular weight of about 1,000 to about 2,500 daltons.

33. A composition according to claim 31 wherein said ULMWH comprise heparin fractions selected from the group consisting of tetrasaccharides, pentasaccharides, hexasaccharides, septasaccharides, octasaccharides and decasaccharides and pharmaceutically acceptable salts thereof.

34. A composition according to claim 31 wherein said ULMWH are N-sulfated.

35. A composition according to claim 31 which comprises about 0.075 to about 0.75 of said ULMWH per kilogram per dos .

- 39 -

36. A liquid or fluid composition according to claim 31 which contains about 1.0 to about 20.0 mg of said ULMWH per ml of composition.

37. A composition according to claim 31 wherein said ULMWH have substantially no anticoagulant activity.

38. A composition according to claim 31 which comprises a solution of said ULMWH in an aqueous, pharmaceutically acceptable inhalant vehicle.

39. A composition according to claim 38 wherein said vehicle is isotonic saline or bacteriostatic water.

40. A composition according to claim 38 which is suitable for administration by means of a pump or squeeze-actuated nebulizer.

41. A composition according to claim 38 which further includes an aerosol propellant and is suitable for administration via a metered aerosol dose inhaler.

42. A composition according to claim 31 which comprises a powdered preparation of said ULMWH intermixed with an inert powder acceptable for intrabronchial administration.

43. A composition according to claim 42 wherein said inert powder is lactose.

44. A composition according to claim 42 which is administered via an aerosol dispenser.

45. A composition according to claim 42 which is administered from a breakable capsule.

46. A composition according to claim 31 which comprises a pharmaceutically acceptable oral vehicle.

- 40 -

47. A composition according to claim 46 which is an oral dosage form selected from the group consisting of tablets, capsules, caplets, gelcaps, pills, liquid solutions, suspensions or elixirs, powders, lozenges, micronized particles and osmotic delivery systems.

48. A composition according to claim 47 which further comprises excipients, vehicles, fillers, binders, disintegrants, solvents, solubilizing agents, sweeteners or coloring agents.

49. A composition according to claim 31 which comprises a pharmaceutically acceptable parenteral vehicle.

50. A composition according to claim 49 wherein said vehicle comprises an isotonic saline solution or a dextrose solution.

51. A composition according to claim 49 which is suitable for intravenous administration.

52. A composition according to claim 49 which is suitable for intramuscular administration.

53. A composition according to claim 31 which comprises a pharmaceutically acceptable topical vehicle.

54. A composition according to claim 31 which comprises a pharmaceutically acceptable intranasal vehicle.

55. A composition according to claim 31 which comprises a pharmaceutically acceptable intraocular vehicle.

- 41 -

56. A method of treating a mammalian patient suffering from or prone to a condition characterized by late phase allergic reactions, airway hyperresponsiveness or inflammatory reactions, said method comprising the administration to the patient of a pharmaceutical composition containing about 0.005 to about 1.0 mg of sulfated polysaccharides per kilogram of patient body weight in each dose, said sulfated polysaccharides having an average molecular weight of about 1,000 to about 3,000 daltons.


57. A method according to claim 56 wherein said sulfated polysaccharides are derived from glycosaminoglycans or mucopolysaccharides.

58. A method according to claim 57 wherein said sulfated polysaccharides are derived from heparin, heparan sulfate, dermatan sulfate, chondroitin sulfate or pentosan polysulfate.

59. A method according to claim 56 wherein said sulfated polysaccharides comprise tetrasaccharides, pentasaccharides, hexasaccharides, septasaccharides, octasaccharides and decasaccharides and pharmaceutically acceptable salts thereof.

60. A method according to claim 59 wherein said sulfated polysaccharides comprise tetrasaccharides.

1/28

FIG. 1

2/28

FIG. 2A





FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7





FIG. 8



9/28

FIG. 9






FIG. 10






12/28

FIG. 12

FRU-70 (1mg/kg PRE-ANTIGEN)







FIG. 14

FRU-70 (0.5 mg/kg POST ANTIGEN)







FIG. 16



17/28

FIG. 17






FIG. 18

HEXASACCHARIDE 0.0625 mg/kg POST ANTIGEN



19/28


FIG. 19



20 / 28

FIG. 20


TETRASACCHARIDE (0.062 mg/kg)



21/28

FIG. 21

0.25 mg/kg POST ANTIGEN (n=4)








FIG. 23




24/28

FIG. 24




25 / 28

FIG. 25




26/28

FIG. 26




27 / 28

FIG. 27



28/28



INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/16111**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(6) :A61K 9/12

US CL :424/45

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/45

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 94/18988 A2 (VON ARNIM) 01 September 1994, see entire document.	1-60

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

21 SEPTEMBER 1998

Date of mailing of the international search report

19 OCT 1998

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

Barthel Lawrence Jr
Raj Bawa, Ph.D.

Telephone No. (703) 308-1235